

Mycoinsecticide - Organic Formulation of BotaniGard

Mycotrol mycoinsecticides are effective biological insecticides that are used to control whitefly, thrips, aphids, and many other insects. They are based on the highly successful fungus, Beauveria bassiana strain GHA, and control the most troublesome crop pests - even resistant strains such as Q-Biotype whitefly. Mycotrol ESO and WPO are versatile enough to use as stand alones, as tank mix partners, or in rotation with other insecticides.



Mycotrol Benefits

- Superior formulation highest spore concentration in a Beauveria-containing insecticide
- · Insect control that equals or exceeds chemical insecticides
- Unique mode of action spores infect directly through the insect's cuticle
- · Excellent tool for resistance management
- · Cost effective controls the early stages of insects, preventing population flare-ups and the expense of rescue treatments
- Excellent safety profile large and diverse crop list and multiple use sites; reduced risk to workers and the environment
- · Compatible for tank-mixes with many pesticides
- Exempt from residue tolerance and has a 0-day pre-harvest interval
- 4-hour REI







Untreated whitefly nymphs







Treated whitefly nymphs

To find out more about the BioWorks family of products, please contact us at 800-877-9443 or visit www.bioworksinc.com.

Unique Mode of Action

Mycotrol's mode of action is unlike any conventional insecticide. The applied spores infect directly through the outside of the insect's cuticle (the skin). Spores adhering to the host will germinate and produce enzymes that attack and dissolve the cuticle, allowing them to penetrate the skin and grow into the insect's body. As the insect dies, it will change color to pink or brown, and eventually the entire body cavity is filled with fungal mass. The most common visible indication of insect death is a discoloration of the immatures. It is not necessary for a white fungal growth to be visible to know the product is working; the insect is killed before this happens.









Size: 1lb.



How You Grow Matters™

100 Rawson Road | Suite 205 Victor, NY 14564 P 800 877 9443 F 800 903 2377 www.bioworksinc.com

Mycotrol[®]

Application

Mycotrol controls a wide variety of insects in greenhouse, field and nursery crops such as whiteflies, thrips, aphids, psyllids, mealybugs, plant bugs and weevils. It has excellent plant safety on a broad range of crops. Mycotrol is registered for an extensive number of ornamentals, herbs and spices, vegetables, fruits and berries. There is no limit to the number of times that Mycotrol can be applied in a season. A weekly application program can prevent insect population explosions and provide a level of control equal to or better than conventional chemical programs.



Mycotrol is Available in Two Formulations:		
Formulation	Pest	Rate
Mycotrol ESO Liquid emulsifiable suspension (WSDA Approved)	Whiteflies, mealybugs, aphids Thrips Other labeled insects	½ qt. to 1 qt./100 gallons 1 qt. to 2 qt./100 gallons ½ qt. to 2 qt./100 gallons*
Mycotrol WPO Wettable powder (WSDA Approved)	Whiteflies, mealybugs, aphids Thrips Other labeled insects Interiorscape bed rates	½ lb to 1 lb/100 gallons 1 lb to 2 lb/100 gallons ½ lb to 2 lb/100 gallons* ¼ to 1 tsp./gal./10 sq.ft.

^{*}Rate depends on insect population and foliage density

Application Considerations:

- Early scouting and detection followed by application of Mycotrol when insect numbers are low result in most effective control.
 Apply at first appearance of insect pests.
- Use a program of Mycotrol applied at 7-day intervals to keep your crop clean and prevent insect population explosions. If applied late or insects get out of hand, reduce the interval to 3-5 days, making at least 3 applications, and use a quick knockdown insecticide to help you regain control.
- Mycotrol is a contact insecticide and thorough coverage is required to achieve control.
- · Mycotrol is safe for use with many beneficial insects.
- Shake the container vigorously and thoroughly to suspend spores.
- Insects are killed long before the appearance of white fungal growth on their surface. The presence of "fuzzy" insect bodies is not required to show product effectiveness.

Note: Use caution when making applications to open blooms, especially on ornamental varieties known to be sensitive. Test on a small group of plants for effects on open blooms before applying to the whole crop. Always read and follow label directions.





100 Rawson Road | Suite 205 Victor, NY 14564 P 800 877 9443 F 800 903 2377 www.bioworksinc.com

Please refer to product labels for complete application details. Additional technical information is available from either our website: www.bioworksinc.com, or your BioWorks sales representative. Always read and follow label directions.

Copyright © 2016 BioWorks Inc., Mycotrol ESO and WPO is a registered trademark of LAM International Corp, Butte, MT