



Metalosate®

Approved for **ORGANIC** Production

Zinc

Amino Acid Chelate Soluble Powder Foliar



Meets NOP Standards

Net Weight: 4.4 pounds (2.0 kg)



08721-AL-KG0002

v02/2009

Lot #:



08000000

METALOSATE® ZINC

Amino Acid Chelate Soluble Powder Foliar

GUARANTEED ANALYSIS (w/w):

Zinc (Zn) 19.3%
19.3% Chelated Zinc

Derived from Zinc Amino Acid Chelate

GENERAL INFORMATION:

Amino acid chelated Metalosate® Zinc soluble powder is designed for foliar application on plants to prevent or correct nutrient deficiencies that may limit crop growth and yields. It is water soluble and nontoxic to plants when applied as directed. For best results, apply Metalosate® Zinc soluble powder according to recommendations based on plant tissue or soil analysis.

DIRECTIONS:

Apply by a spray method and in an adequate amount of water that will provide complete coverage of the plants. Use of a non-ionic wetting agent may improve spray coverage of certain hard to wet plants.

Metalosate® Zinc soluble powder can be included in a regular spray program on crops. Consult with an Albion® representative on compatibility with other spray materials.

The rate of application will depend on the crop, stage of growth, and severity of deficiency. The maximum recommended rates are for mature, full-sized plants. Reduce the rates proportionately when spraying smaller plants.

Metalosate® Zinc soluble powder may be used on legumes, grain crops, root crops, cucurbits, cole crops, leafy vegetables, woody and herbaceous ornamentals, deciduous fruits, vine crops, tropical and subtropical fruits, and many other crops.

Approved for
ORGANIC
Production

RECOMMENDATIONS:

Field Crops and Vegetables: Apply 7 to 14 ounces (200 to 400 grams) per acre during periods of rapid growth or nutritional stress. The application may be repeated 2 or more times through the growing season.

Tree Crops: Make an application of 7 to 21 ounces (200 to 600 grams) per acre after the beginning of active growth. The application may be repeated at 2 to 4 week intervals through the growing season. On deciduous trees, 7 to 14 ounces (200 to 400 grams) per acre may be applied in the oil sprays at the delayed dormant stage; post-harvest applications of 14 to 21 ounces (400 to 600 grams) per acre may be made while active, green leaves remain on the trees.

Grapes and Berries: Make an application of 7 to 14 ounces (200 to 400 grams) per acre after active growth begins. The application may be repeated at intervals of one week or more through the vegetative-growth period.

LIMITED WARRANTY:

Seller's and Manufacturer's obligation is limited to replacement of defective product. **NEITHER SELLER NOR MANUFACTURER SHALL BE LIABLE FOR ANY INJURY, LOSS, OR DAMAGE DIRECTLY OR CONSEQUENTLY ARISING OUT OF THE MISUSE OR INABILITY TO USE THE PRODUCT. ALL OTHER WARRANTIES EXCEPT IN WRITING FROM THE MANUFACTURER, WHETHER EXPRESSED OR IMPLIED ARE HEREBY DISCLAIMED.**

General information regarding the contents and levels of metals in this product is available by calling 1-866-CHELATE (1-866-243-5283).

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>

Albion, the Albion logo, and Metalosate are trademarks of Albion Laboratories, Inc.

Reseal package between mixings to maintain freshness.

This product has been
verified to comply with 7 CFR, Part 205
United States Department of Agriculture
National Organic Program

U.S. Patent Pending F62

Manufactured by:

Albion Laboratories, Inc.
101 North Main Street
Clearfield, Utah 84015 U.S.A.

ALv02/2009

