



BorreGro® HA-2 (LIQUID)

Humic Acid



BorreGro® HA-2 promotes the uptake of micronutrients.

BorreGro® HA-2 may be applied alone or with liquid fertilizers with a pH of 7.0 or higher.

Compatibility can be simply checked by:

- Mixing each product in a clear container in the same proportion to be used in the field
- Agitating vigorously for at least 30 seconds
- Visually observing for precipitate

Always consult the Material Safety Data Sheet for instructions on proper handling.

Notice: The statements made on this label are believed to be true and accurate but because of conditions of use which are beyond our control LignoTech USA does not authorize any agent or representative to make any warranty, guarantee or representation, expressed or implied, concerning this material or the use thereof, except in conformity with the statements on this label. Neither LignoTech USA nor the seller shall not be responsible in any manner for any property damage or personal injury resulting from the use of this material not in accordance with directions. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions.

GUARANTEED ANALYSIS

15% Humic Acid derived from Leonardite
(BaCl₂ Method)

“ALSO CONTAINS NON-PLANT FOOD
INGREDIENTS”

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin and clothing. In case of contact, immediately flush with water.

Contact your LignoTech Sales Representative for additional information.

**Manufactured in U.S.A. by
LignoTech USA, Inc.**
100 Grand Avenue
Rothschild, WI 54474
Tel: (715)-359-6544
Fax: (715)-355-3674

Net Contents: 275 gal (1040 liters)
Wt. Per gallon: 9.10 lbs. @ 70°F (1.09 kg/l)
Net Weight: 2502 lbs (1137 kgs)

GUIDELINES FOR USE:

BorreGro® HA-2 can be applied to all crops including fruits, vegetables, trees, vines, flowers, ornamentals, turf, grain and other row crops.

Application rates and frequency are dependent upon soil and climate conditions present. Typical application rates range from ½ to 10 pounds per crop per acre per season.

BorreGro® HA-2 may be applied through conventional methods but not limited to foliar spray, soil application, drip irrigation, water run, shanking in or hydroponically.

Follow the labeled rates for nutrients and/or consult with your local PCA.

Lot Number: _____

Information regarding the contents and levels of metals in this product is available on the internet at
<http://www.aapfco.org/metals.htm>

CO, ME, MI, NV, NY, ND, PA, WA

