

Supplemental Labeling



Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Entrust® SC

EPA Reg. No. 62719-621

Tropical Tree Fruits¹

¹Acerola, atemoya, avocado, biriba, black sapote, canistel, cherimoya, custard apple, feijoa, guava, ilama, jaboticaba, longan, lychee, mamey sapote, mango, papaya, passionfruit, pitaya (dragon fruit), pulasan, rambutan, sapodilla, soursop, Spanish lime, star apple, starfruit, sugar apple, ti leaves, wax jambu, white sapote

This label expires and must not be distributed or used in accordance with this supplemental registration after March 26, 2024.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Entrust® SC insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Entrust SC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Entrust SC.

Directions for Use

Pests and Application Rates:

| Pests | Entrust SC (fl oz/acre) |
|--|-------------------------|
| European grapevine moth katydids lepidopterous larvae avocado leafroller citrus peelminer cutworms fruit tree leafroller naval orange worm orange tortrix western tussock moth light brown apple moth thrips ¹ | 4 - 10 |

¹Control of thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants section under Mixing Directions.

Application Timing: Treat when pests appear or in accordance with local economic thresholds. Consult your Corteva Agriscience representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: The amount of Entrust SC applied per acre will depend upon tree size and pest pressure. Use a lower rate in the rate range for light infestations and/or small trees and a higher rate in the rate range for heavy infestations and/or large trees.

Resistance Management: Do not make more than two consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after two consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your local Corteva Agriscience representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Restrictions:

- **Preharvest Intervals:** Do not apply within 1 day of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 7 days apart.
- Do not apply more than a total of 29 fl oz of Entrust SC (0.45 lb ai spinosad) per acre per crop for all methods of application, including foliar, soil and seed treatment uses.
- **Maximum Number of Applications:** In order to prevent or delay resistance development in thrips, do not apply Entrust SC more than two times per year.
- For resistance management purposes, do not apply to tropical tree fruits grown in nurseries or in greenhouses.

™®Trademarks of Corteva Agriscience and its affiliated companies

R399-038
EPA accepted 04/09/2021
Initial printing or Replaces R399-036